



Altosid[®]

LIQUID LARVICIDE

MOSQUITO GROWTH REGULATOR

**PREVENTS EMERGENCE OF ADULT FLOODWATER MOSQUITOES
FOR CONTROL OF MOSQUITO LARVAE INCLUDING ULV APPLICATION**

SPECIMEN LABEL

ACTIVE INGREDIENT:

(S)-Methoprene (CAS #65733-16-6).....5%
OTHER INGREDIENTS:95%
TOTAL: 100%

Formulation contains 0.43 lb/gal (51.3 g/liter) active ingredient
EPA REG. NO. 2724-392 EPA EST. NO. 2724-TX-1

KEEP OUT OF REACH OF CHILDREN
CAUTION
SEE ADDITIONAL PRECAUTIONARY STATEMENTS

BECAUSE OF THE UNIQUE MODE OF ACTION OF **A.L.L.**, SUCCESSFUL USE REQUIRES FAMILIARITY WITH SPECIAL TECHNIQUES FOR APPLICATION TIMING AND TREATMENT EVALUATION. SEE **GUIDE TO PRODUCT APPLICATION** OR CONSULT LOCAL MOSQUITO ABATEMENT AGENCY.

PRECAUTIONARY STATEMENTS – HAZARDS TO HUMANS – AND DOMESTIC ANIMALS
CAUTION

Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

FIRST AID

If in eyes • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.

(continued)

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-248-7763 for emergency medical treatment information.

ENVIRONMENTAL HAZARDS

Do not contaminate water when disposing of equipment washwaters or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

CHEMIGATION

Refer to supplemental labeling entitled **Guide to Product Application** for use directions for chemigation. Do not apply this product through any irrigation system unless the supplemental labeling on chemigation is followed.

MIXING AND HANDLING INSTRUCTIONS

1. **SHAKE WELL BEFORE USING.** Zoëcon[®] Altosid[®] Liquid Larvicide Mosquito Growth Regulator (A.L.L.) may separate on standing and must be thoroughly agitated prior to dilution.
2. Do not mix with oil; use clean equipment.
3. Partially fill spray tank with water; then add the labeled amount of **A.L.L.**, agitate and complete filling. Mild agitation during application is desirable.
4. Use spray solution within 48 hours; always agitate before spraying.

APPLICATION INSTRUCTIONS

Introduction: **A.L.L.** must be applied to 2nd, 3rd, or 4th larval instars of floodwater mosquitoes to prevent adult emergence. Treated larvae continue normal development to the pupal stage where they die. This insect growth regulator **has no effect when applied to pupae or adult mosquitoes.** **A.L.L.** has sufficient field life to be effective at appropriate rates when applied to larval stages under varying field conditions. For further information, see **Guide to Product Application.**

METHODS OF APPLICATION

Aerial: Use the prescribed amount of **A.L.L.** listed below in sufficient water to give complete coverage. One-half to 5 gallons of spray solution per acre is usually satisfactory. Do not apply when weather conditions favor drift from areas treated.

Ground: Determine the average spray volume used per acre by individual operators and/or specific equipment. Mix **A.L.L.** in the appropriate volume of water to give the rate per acre as indicated below.

For Control of Mosquito Larvae Using ULV Application:

For ground application to terrestrial sites, apply at the rate of 3-4 ounces of product per acre to water-holding containers and other small bodies of water that breed mosquitoes. Use equipment capable of applying a fine mist or ULV. Follow equipment manufacturer's recommendations when making applications. Direct spray applications to sites where mosquitoes breed. These sites include: tires and tire piles, potted plants, tree holes, garbage bins, cans, birdbaths, rain barrels, and other water-holding containers and small bodies of water.

APPLICATION RATE

Apply 3 to 4 fl oz of **A.L.L.** per acre (219 to 293 ml/hectare) in water as directed. **For Control of Mosquito Larvae Using ULV Application:** Apply at 3-4 ounces per acre. Reapply as breeding sites become reinfested or when monitoring indicates an increase in adult populations.

APPLICATION SITES

Pastures: **A.L.L.** may be applied after each flooding without removal of grazing livestock.

Rice: **A.L.L.** must be applied to 2nd, 3rd, and/or 4th instar larvae of mosquitoes found in rice, usually within 4 days after flooding. **A.L.L.** treatment may be repeated with each flooding.

Intermittently Flooded Noncrop Areas: **A.L.L.** may be applied as directed above when flooding may result in floodwater mosquito hatch. Typical sites include: freshwater swamps and marshes, salt marshes, woodland pools and meadows, dredging spoil sites, drainage areas, waste treatment and settling ponds, ditches and other natural and manmade depressions.

Crop Areas: **A.L.L.** may be applied to irrigated croplands after flooding to control mosquito emergence. Examples of such sites are: vineyards, rice fields (including wild rice), date palm orchards, fruit and nut orchards, and berry fields and bogs. Irrigated pastures may be treated after each flooding without the removal of grazing livestock.

Dense Vegetation or Canopy Areas: Apply an **A.L.L.** sand or BIODAC® mixture using standard granular dispersal equipment. For detailed preparation instructions, refer to **Guide to Product Application**.

TANK MIXING INSTRUCTIONS

A.L.L. may be tank mixed with liquid *Bacillus thuringiensis*, variety *israeliensis* (*B.t.i.*) formulations. The ratio of *B.t.i.* to **A.L.L.** ranges from 6:1 to 12.5:1 (volume/volume). For example, to prepare a 12.5:1 tank mix, add 1 gallon of **A.L.L.** to 12.5 gallons of *B.t.i.* This tank mix can be applied to sites listed above at rates of 2–16 fluid ounces/acre (0.15-1.2 liters/hectare).

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. **Storage:** Store in cool place. Store product away from other pesticides, food, and feed. In case of leakage or spill, soak up with sand or another absorbent material. **Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **Container Disposal:** Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying. Triple rinse as follows. Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

For information, call **1-800-248-7763** or visit our Web site: **www.altosid.com**

Wellmark International
1501 East Woodfield Road 200W
Schaumburg, Illinois 60173



Altosid, Zoëcon and the Zoëcon logo are registered trademarks of Wellmark International.
©2006-2011 Wellmark International.

October, 2011
Schaumburg, IL