



Photo courtesy of Ohio State Weed Lab Archive, The Ohio State University, Bugwood.org.

Nufarm Prosedge®

SELECTIVE HERBICIDE

Stop Nutsedge in its Tracks with Nufarm Prosedge™.

Yellow nutsedge can be one of the most troublesome weeds found in lawns and other turfgrass: unlike most weeds, it's not controlled with applications of traditional grass or broadleaf herbicides. Nufarm Prosedge™ offers an alternative for selective nutsedge control in turf and around woody ornamentals in landscaped areas.

FORMULATED FOR NUTSEGE CONTROL

- Excellent control of yellow and purple nutsedge and horsetail
- Kyllinga suppression
- Easy-to-use water-soluble formulation
- Rainfast within 4 hours of spraying
- Gentle to many turfgrass species

EPA REG. NO. 228-702

ACTIVE INGR. halosulfuron-methyl (75.0%)

FORMULATION water-dispersible granule

CHEM. FAMILY sulfonyleurea

HRAC NO. 2

SIGNAL WORD caution

RESTRICTED no

PACKAGE SIZE 10 x 1.33 oz bottle,
30 x 0.9 gm water-soluble
packet

 RESIDENTIAL

 COMMERCIAL

 TURF

 LANDSCAPE ORNAMENTALS

 GOLF TURF

 POST-EMERGE



KEY USES

Turfgrasses (*established lawns, ornamental turfgrass, landscaped areas, commercial and residential turfgrass*)

Non-crop sites (*including airports, cemeteries, fallow areas, golf courses, landscaped areas, public recreation areas, rights-of-way, etc.*)

KEY WEEDS CONTROLLED

(*See product label for complete list*)

Horsetail

Purple nutsedge

Yellow nutsedge

APPLICATION RECOMMENDATIONS

FOR POST-EMERGENCE CONTROL OF PURPLE OR YELLOW NUTSEDGE

For post-emergence control of purple or yellow nutsedge found in established turfgrass, apply 2/3 to 1 1/3 ounces by weight per acre after nutsedge has reached the 3 to 8 leaf stage of growth.

If a second treatment is required, apply 6 to 10 weeks after the initial treatment. As a sequential treatment, when new purple or yellow nutsedge plants have reached the 3 to 8 leaf stage of growth, apply 2/3 to 1 1/3 ounces by weight per acre.

Use 0.25 to 0.5% nonionic surfactant, or 1-2 quarts per 100 gallons broadcast, or 2 teaspoons (1/3 fl oz) per gallon spot treatment.

When applied as directed, the following established turfgrasses are tolerant to application of this product:

Cool-season grasses: creeping bentgrass, fine fescue, Kentucky bluegrass, perennial ryegrass, tall fescue

Warm-season grasses: bahiagrass, bermudagrass, centipedegrass, kikuyugrass, seashore paspalum, St. Augustinegrass, zoysiagrass

FOR POST-EMERGENCE CONTROL OF HORSETAIL

For post-emergence control of horsetail (*Equisetum arvense*) in roadsides, rights-of-way, tank farms, lumber yards, fuel storage areas, fallow land and fence rows, apply 2 2/3 ounces per acre or 0.06 ounces (1.8 grams) per 1,000 square feet after horsetail has leafed out.

See label for complete application rates and recommendations.

